

EFFECT OF Î±-TOCOPHEROL AND ASCORBIC ACIDS ON PERFORMANCE AND BLOOD IMMUNITY PROFILE OF MALE NATIVE MUSCOVY DUCK

Publons ID	36331702
Wos ID	WOS:000417320100005
Doi	10.14710/jitaa.41.3.145-152
Title	EFFECT OF Î±-TOCOPHEROL AND ASCORBIC ACIDS ON PERFORMANCE AND BLOOD IMMUNITY PROFILE OF MALE NATIVE MUSCOVY DUCK
First Author	Tugiyanti, E.; Yuwanta, T.; Zuprizal;
Last Author	Ismoyowati
Authors	Tugiyanti, E; Yuwanta, T; Zuprizal; Rusman; Ismoyowati;
Publish Date	SEP 2016
Journal Name	JOURNAL OF THE INDONESIAN TROPICAL ANIMAL AGRICULTURE
Citation	2
Abstract	<p>The purpose of the research was to assess the effect of alpha-tocopherol and ascorbic acids on the performance and immunity blood profile of male native Muscovy duck. The materials used were 84 male Muscovy ducks at 9 weeks old. The research used completely randomized design with 7 treatments. The treatments were E0C0 : basal diet without alpha-tocopherol and ascorbic acids, E400: basal diet+ alpha-tocopherol 400IU, E600: basal diet+alpha-tocopherol 600IU, C400: basal diet+ascorbic acid 400mg/kg, C600: basal diet+ascorbic acid 600mg/kg, E200C200: basal diet+alpha-tocopherol 200IU+ ascorbic acid 200mg/kg, and E300C300: basal diet+alpha-tocopherol 300IU+ ascorbic acid 300mg/kg. Each treatment was repeated 4 times and each replication consisted of 3 ducks. The observed variables were body weight, feed consumption, feed conversion ratio, packed cell volume (PCV), total of plasma protein (TPP), leucocyte count, basophil, heterophil, eosinophil, monocyte, lymphocytes count and heterophil/lymphocyte (H/L) ratio of Muscovy duck. The result indicated that E0C0, E400, E600, C400, C600, E200C200 and E300C300 had no significant difference in body weight, feed intake and feed conversion. There were significant effects on PCV, leucocyte count, percentage of heterophil and lymphocytes, but had no significant effect on eosinophil, monocyte and the H/L ratio. The C400 resulted a higher PCV count, percentage of heterophil, and H/L ratio. The C 600 produced the highest leucocyte count. In conclusion, the supplementation of ascorbic acid at 400 - 600 mg/kg feed ascorbic acid could improve the immune profile, but could not improve the performance of Muscovy duck.</p>
Publish Type	Journal
Publish Year	2016
Page Begin	145
Page End	152
Issn	2087-8273
Eissn	2460-6278
Url	https://www.webofscience.com/wos/woscc/full-record/WOS:000417320100005
Author	Dr Ir ELLY TUGIYANTI, M.P