

Paradoxical sleep deprivation changes testicular malondialdehyde and caspase-3 expression in male rats

Publons ID	44514893
Wos ID	WOS:000376161700003
Doi	10.18051/UnivMed.2015.v34.87-95
Title	Paradoxical sleep deprivation changes testicular malondialdehyde and caspase-3 expression in male rats
First Author	
Last Author	
Authors	Arjadi, F; Partadireja, G; Maurits, LS; Pangestu, M;
Publish Date	MAY-AUG 2015
Journal Name	UNIVERSA MEDICINA
Citation	
Abstract	<p>a:8:{i:0;s:10:"BACKGROUND";i:1;s:578:"Sleep deprivation is a significant problem among adult men and is considered as a risk factor for several diseases. Paradoxical sleep deprivation (PSD) induces Leydig cell apoptosis through elevation of corticosterone, with testicular malondialdehyde (MDA) and Leydig cell caspase-3 expression as parameters. The aim of this study was to observe testicular MDA level and caspase-3 expression treated with paradoxical sleep deprivation (PSD), immobilization, and footshock stress and to determine the stress model with a significant effect in white male rats (<i>Rattus norvegicus</i>).";i:2;s:7:"METHODS";i:3;s:458:"This experimental randomized study of posttest only with control group design was conducted on 24 white male Wistar strain rats, randomly allocated into four treatment groups, i.e. control (K1) without any stress treatment, PSD (KII), immobilization (KIII), and footshock stress (KIV). Treatments were given for 25 days to produce chronic stress. Testicular MDA concentration was examined by the ELISA method while caspase-3 was examined by the TUNEL method.";i:4;s:7:"RESULTS";i:5;s:472:"Mean testicular MDA concentration with one-way ANOVA test showed differences in means between the groups ($p=0.000$) and post hoc Tukey-HSD test showed significant results between PSD stress group versus control, immobilization and footshock stress groups. One-way ANOVA test showed a significant difference in caspase-3 expression in at least two treatment groups ($p=0.008$) and post-hoc Tuckey-LSD test showed significant differences between controls and all stress groups.";i:6;s:10:"CONCLUSION";i:7;s:131:"Sleep deprivation is a type of stress inducing changes in testicular MDA concentration and caspase-3 expression in male rat testes.";}</p>
Publish Type	Journal
Publish Year	2015
Page Begin	87
Page End	95
Issn	1907-3062
Eissn	2407-2230
Url	https://www.webofscience.com/wos/woscc/full-record/WOS:000376161700003
Author	Dr Dr FITRANTO ARJADI, S.Ked, M.Kes