

A First-Principles Study of the Adsorption of H₂O on Ru- and Mo-Alloyed Pt(111) Surfaces

Publons ID	31478854
Wos ID	WOS:000515822800036
Doi	10.1007/s11664-020-07976-4
Title	A First-Principles Study of the Adsorption of H ₂ O on Ru- and Mo-Alloyed Pt(111) Surfaces
First Author	Cahyanto, Wahyu Tri; Zulaehah, Siti; Abdullatif, Farzand;
Last Author	Kasai, Hideaki
Authors	Cahyanto, WT; Zulaehah, S; Abdullatif, F; Widanarto, W; Effendi, M; Kasai, H;
Publish Date	APR 2020
Journal Name	JOURNAL OF ELECTRONIC MATERIALS
Citation	6
Abstract	<p>A study on the molecular adsorption of a water (H₂O) monomer on Pt alloy surfaces with binary Pt-Ru and Pt-Mo and ternary Pt-Ru-Mo surface models was conducted. Some calculations of the interaction between a H₂O monomer and the Pt surface were also done for reference. This study is based on density functional theory (DFT) with periodic models and aims to understand the H₂O adsorption mechanism. In Pt, Pt-Ru, and Pt-Mo surfaces, H₂O preferably adsorbs via an oxygen atom in a flat configuration at the top sites of Pt, Ru, Mo, respectively. However, on the ternary Pt-Ru-Mo surface, the adsorption configuration has the most inclined H₂O structure relative to the surface. The results showed that the binding energy of H₂O/Pt-Ru-Mo > H₂O/Pt-Mo > H₂O/Pt-Ru > H₂O/Pt. The adsorption mechanism was then clarified by charge transfer and natural bonding. The charge transfer from the surface to the adsorbate is observed in all models, with the greatest charge transfer occurring on the surface of Pt doped with two Mo atoms. This is probably due to the fact that oxygen can attract the most charge on Mo, because the difference in electronegativity is greatest. The calculation results also show that Ru is the most hydrophilic metal for oxygen. However, since the adsorption structure is parallel to the surface, hydrogen (H) is also more sensitive to receive the charge. Subsequently, the most acceptable reason for the most stable adsorption for H₂O/Pt-Ru-Mo is that the inclined structure yields the most orbitals hybridization at the H₂O's highest occupied molecular orbital (HOMO). This drives the interaction by forming bonding states at the lowest energy and anti-bonding states at the highest energy.</p>
Publish Type	Journal
Publish Year	2020
Page Begin	2642
Page End	2650
Issn	0361-5235
Eissn	1543-186X
Url	https://www.webofscience.com/wos/woscc/full-record/WOS:000515822800036
Author	WAHYU TRI CAHYANTO, S.Si, M.Si, Ph.D