Chiral separation of vinpocetine using cyclodextrin-modified micellar electrokinetic chromatography

Publons ID	4140054
Wos ID	WOS:000300664000009
Doi	10.1002/chir.21990
Title	Chiral separation of vinpocetine using cyclodextrin-modified micellar electrokinetic chromatography
First Author	
Last Author	
Authors	Ibrahim, WAW; Abd Wahib, SM; Hermawan, D; Sanagi, MM; Aboul-Enein, HY;
Publish Date	MAR 2012
Journal Name	CHIRALITY
Citation	15
Abstract	A cyclodextrin-modified micellar electrokinetic chromatography (CD-MEKC) technique has been developed for enantioseparation of vinpocetine using an inexpensive 2-hydroxypropyl-beta-CD (HP-beta-CD) as the chiral selector (CS). The best chiral separation was achieved using 40 mM HP-beta-CD as the CS in 50 mM phosphate buffer (pH 7.0) consisting of 40 mM sodium dodecyl sulfate (SDS) at a separation temperature and separation voltage of 25 degrees C and 25 kV, respectively. To the author's best knowledge, this is the first CD-MEKC study able to successfully separate the four stereoisomer of vinpocetine in separation time of 9.5 min and resolution of 1.043.87. Chirality, 2012. (C) 2012 Wiley Periodicals, Inc.
Publish Type	Journal
Publish Year	2012
Page Begin	252
Page End	254
lssn	0899-0042
Eissn	1520-636X
Url	https://www.webofscience.com/wos/woscc/full-record/WOS:000300664000009
Author	DADAN HERMAWAN