Synthesis of Ag3PO4 using Hydrophylic Polymer and Their Photocatalytic Activities under Visible Light Irradiation

Doi 10	VOS:000401834200009 0.9767/bcrec.12.2.767.206-211
Doi 10	
S.	11 U/6 / bcroc 12 2 / 6 / 206 211
- Sy	
Title Lig	Synthesis of Ag3PO4 using Hydrophylic Polymer and Their Photocatalytic Activities under Visible ight Irradiation
First Author	
Last Author	
Authors Su	Sulaeman, U; Liu, B; Yin, S; Sato, T;
Publish Date	NUG 2017
Journal Name	BULLETIN OF CHEMICAL REACTION ENGINEERING AND CATALYSIS
Citation 4	
Abstract po Fie us the of Co	The highly active Ag3PO4 photocatalysts were successfully synthesized using the hydrophylic olymer of PVA (polyvinyl alcohol), PEG (polyethylene glycol) and PVP (polyvinyl pyrrolidone). The products were characterized using X-ray diffraction (XRD), Diffuse reflection spectroscopy (DRS), Tield emission scanning electron microscope (FE-SEM), Brunauer-Emmett-Teller (BET) specific urface area, and Xray photoelectron spectroscopy (XPS). Photocatalytic activities were evaluated using decomposition of Rhodamine B (RhB) under visible light irradiation. The results showed that the PVA, PEG, and PVP increased the specific surface area and enhanced the photocatalytic activity of Ag3PO4. The highest photocatalytic activity could be observed in Ag3PO4 synthesized with PVA, hainly due to an increase in electron excitation induced by PVA chemically adsorbed on the surface. Copyright (C) 2017 BCREC Group. All rights reserved
Publish Type	ournal
Publish Year	017
Page Begin	06
Page End 21	11
Issn 19	978-2993
Eissn	
Url htt	ttps://www.webofscience.com/wos/woscc/full-record/WOS:000401834200009
Author UY	JYI SULAEMAN, S.Si, M.Si, PhD