

**POLA PENYERAPAN KUNING TELUR DAN PERKEMBANGAN ORGANOGENESIS
PADA STADIA AWAL LARVA IKAN SENGGARINGAN (*Mystus nigriceps*)**

Title	POLA PENYERAPAN KUNING TELUR DAN PERKEMBANGAN ORGANOGENESIS PADA STADIA AWAL LARVA IKAN SENGGARINGAN (<i>Mystus nigriceps</i>)
Author Order	1 of 2
Accreditation	
Abstract	The observation was conducted to study the yolk absorption activity andmorphological development on the initial state of senggaringan (catfish) fishlarvae. Larvae used for this research were from the artificial spawning ofbroodstock, reared in fiber tank. The result showed that newly hatched larvae has3.02 mm $\text{Lf} \pm 0.14$ total length and 1.1202 mm ³ yolk volume. During yolk absorptionperiod, organogenesis development and length added also occurred. Larvaedevelopment occurred after hatching involved the eye spot (23 hours, 1 minute),mouth opening (25 hours, 21 minutes) and pigmentation (50 hours, 5 minutes),. Organogenesis development and the total length increment improve, following theincrease of yolk absorption rate.
Publisher Name	Fakultas Perikanan dan Kelautan, Universitas Riau
Publish Date	2012-11-21
Publish Year	2009
Doi	DOI: 10.31258/terubuk.37.1.%p
Citation	
Source	Berkala Perikanan Terubuk
Source Issue	Vol 37, No 1 (2009): Februari 2009
Source Page	
Url	https://terubuk.ejournal.unri.ac.id/index.php/JT/article/view/223/217
Author	Dr TAUFIK BUDHI PRAMONO, S.Pi, M.Si