

A MONOGRAPH OF THE GENUS SCHOUTENIA*) Korth. (Tiliaceae)

Title	A MONOGRAPH OF THE GENUS SCHOUTENIA*) Korth. (Tiliaceae)
Author Order	2 of 2
Accreditation	
Abstract	The genus consists of 8 species in Siam, Indochina, the Malay Peninsula, Sumatra, Borneo, Java, and the Lesser Sunda Islands. It is here divided into 2 subgenera, the monotypic Schoutevia with <i>S. cvata</i> Korth. and Chartacalyx with <i>S. accrescens</i> (Mast.) Curtis as type species. Chartacalyx is divided into 2 sections, Chartacalyx and A disci flora with <i>S. kuiistleri</i> King as type spscies. Three species, <i>S. curtisii</i> , <i>corneri</i> , and <i>kostermansii</i> are considered new. <i>S. hypoleuca</i> Pierre is reduced to synonymy of <i>S. ovata</i> Korth. <i>S. peregrines</i> Craib is reduced to a subspecies of <i>S. glomerata</i> King. The varieties of <i>S. accrescens</i> (Mast.) Curtis are deleted. The species is divided into 3 subspecies: <i>accrescens</i> , <i>stellata</i> , and <i>borneensis</i> . A forma <i>lepidota</i> is reeognized for subspecies <i>accrescens</i> . <i>S. buurmanii</i> K. et V. and <i>S. kunstleri</i> King- are considered to be different species.
Publisher Name	Research Center for Biology
Publish Date	2015-11-17
Publish Year	1965
Doi	DOI: 10.14203/reinwardtia.v7i2.972
Citation	
Source	REINWARDTIA
Source Issue	Vol 7, No 2 (1965): vol.7 no.2
Source Page	91-138
Url	http://e-journal.biologi.lipi.go.id/index.php/reinwardtia/article/view/972/826
Author	HARTONO, M.Si