

The Starting Material Concentration Dependence of Ag₃PO₄ Synthesis for Rhodamine B Photodegradation under Visible Light Irradiation

Title	The Starting Material Concentration Dependence of Ag ₃ PO ₄ Synthesis for Rhodamine B Photodegradation under Visible Light Irradiation
Author Order	2 of 2
Accreditation	2
Abstract	Synthesis of Ag ₃ PO ₄ photocatalyst under the varied concentrations of AgNO ₃ and Na ₂ HPO ₄ ·12H ₂ O as starting material has been successfully synthesized using the co-precipitation method. The concentration of AgNO ₃ is 0.1; 0.5; 1.0; and 2.0 M, whereas Na ₂ HPO ₄ ·12H ₂ O is 0.03; 0.17; 0.33; and 0.67 M, respectively. The co-precipitations were carried out under aqueous solution. As-synthesized photocatalysts were examined to degrade Rhodamine B (RhB) under blue light irradiation. The results showed that varying concentrations of starting materials affect the photocatalytic activities, the intensity ratio of [110]/[200] facet plane, and their bandgap energies of Ag ₃ PO ₄ photocatalyst. The highest photocatalytic activity of the sample was obtained by synthesized using the 1.0 M of AgNO ₃ and 0.33 M of Na ₂ HPO ₄ ·12H ₂ O (AP-1.0). This is due to the high [110] facet plane and increased absorption along the visible region of AP-1.0 photocatalyst. Therefore, this result could be a consideration for the improvement of Ag ₃ PO ₄ photocatalyst.
Publisher Name	Syarif Hidayatullah State Islamic University
Publish Date	2020-05-29
Publish Year	2020
Doi	DOI: 10.15408/jkv.v6i1.14837
Citation	
Source	Jurnal Kimia Valensi
Source Issue	Jurnal Kimia VALENSI Volume 6, No. 1, May 2020
Source Page	1-9
Url	https://journal.uinjkt.ac.id/index.php/valensi/article/view/14837/pdf
Author	UYI SULAEMAN, S.Si, M.Si, PhD