Amount of Menstrual Blood and Nutrient Intake with Hemoglobin Level

Title	Amount of Menstrual Blood and Nutrient Intake with Hemoglobin Level
Author Order	2 of 2
Accreditation	2
Abstract	The most influential factor in the anemia is severe menstrual bleeding on each cycle. Heavy menstrual bleeding that can periodically reduce iron stores in the body so that the body has an iron deficiency and lead to anemia. This study aimed to find out the relationship between menstrual blood counts and nutritional intake with hemoglobin (Hb) levels in girls. The study used a cross-sectional design with a sample of 169 young girls in middle school in Banyumas District, Central Java, Indonesia. The amount of menstrual blood was seen using a questionnaire menstrual pictogram. The nutritional intake studied was the intake of protein, fibre, iron, and vitamin C using an FFQ semi-quantitative questionnaire and Hb levels through the insertion of peripheral blood. Data analysis using is chi-square and Fisher exact tests. There was no significant relationship between the amount of menstrual blood with Hb levels (p = 0.54 Cl 95% = 0.36-1.74), protein intake (p = 0.26), fibre intake (p = 0.78) and iron intake (p = 0.44). There was a significant relationship between vitamin C intake and Hb levels (p = 0.03 Cl 95% = 1.04-4.10). There is a significant relationship between vitamin C intake and hemoglobin levels in young girls. $\tilde{A}f\hat{A},\tilde{A},\tilde{A}$
Publisher Name	Department of Public Health, Faculty of Sport Science, Universitas Negeri Semarang
Publish Date	2021-07-24
Publish Year	2021
Doi	DOI: 10.15294/kemas.v17i1.24540
Citation	
Source	KEMAS: Jurnal Kesehatan Masyarakat
Source Issue	Vol 17, No 1 (2021)
Source Page	31-36
Url	https://journal.unnes.ac.id/nju/index.php/kemas/article/downloadSuppFile/24540/5562
Author	HIYA ALFI RAHMAH, S.Gz, Dietisien, M.P.H