

Prediction Models for Decreasing Visual Acuity in Wig Makers

Title	Prediction Models for Decreasing Visual Acuity in Wig Makers
Author Order	3 of 3
Accreditation	1
Abstract	<p>One of the occupational diseases that can arise for workers with high accuracy is a decrease in their visual acuity. Therefore, it is necessary to study the risk factors of decreasing visual acuity in workers with high accuracy, such as in wig makers. This study aimed to examine the correlation between age, working period, lighting intensity, fatigue, and nutritional status with visual acuity, and to observe the main risk factors that can be used as a reference for predicting decreasing visual acuity. This study was an observational study that used a cross-sectional design. The population number for this study was 185 wig makers. A total of 85 samples were selected using simple random sampling. The study was carried out in January to March 2016 on wig makers in Purbalingga. The results showed that variables most closely associated with visual acuity were working period ($p\text{-value} = 0.000 < 0.05$), lighting intensity ($p\text{-value} = 0.000 < 0.05$), and fatigue ($0.013 < 0.05$). Variables that were not related to vision acuity were age ($p\text{-value} = 0.846 > 0.05$) and nutritional status ($p\text{-value} = 0.562 > 0.05$).</p>
Publisher Name	Faculty of Public Health Universitas Indonesia
Publish Date	2019-02-04
Publish Year	2019
Doi	DOI: 10.21109/kesmas.v13i3.1784
Citation	
Source	Jurnal Kesehatan Masyarakat Nasional
Source Issue	Volume 13, Issue 3, February 2019
Source Page	138-143
Url	http://journal.fkm.ui.ac.id/kesmas/article/downloadSuppFile/1784/967
Author	Dr Drs NGADIMAN, M.Kes, M.Kes