

Plantlet Formation and Acclimatization of Sugarcane cv. PS 881 with Different Types and Concentration of Auxin

Title	Plantlet Formation and Acclimatization of Sugarcane cv. PS 881 with Different Types and Concentration of Auxin
Author Order	2 of 4
Accreditation	2
Abstract	<p>This research has been carried out with a view to induce rooting and plantlet formation, followed by acclimatization. Micro shoots of sugarcane cv. PS 881 were cultured on Murashige and Skoog medium supplemented with different types and concentration of auxins for root induction. This research conducted experimentally using a split-plot design. The main plots were three types of auxins, which consisted of IAA, IBA, and NAA. The subplots were auxin concentrations with four levels, i.e. 0 μM, 5 μM, 10 μM, and 15 μM. Significantly faster root emergence time and higher number of roots observed in the Murashige and Skoog basal medium supplemented with 10 μM NAA. The best root length obtained in the Murashige and Skoog basal medium supplemented with NAA 0 μM. Plantlets derived from NAA 10 μM treatment showed the best performance during acclimation with a 100% survival rate. NAA at a concentration of 10 μM considered to be the best treatment in plantlet formation and acclimatization of sugarcane cv. PS 881. This study showed that the use of MS medium with 10 μM NAA is able to increase the growth of PS 881 sugarcane plantlets. The results of this study can increase the availability of high quality seedlings and increase national sugar production.</p>
Publisher Name	Department of Biology, Faculty of Mathematics and Sciences, Semarang State University . Ro
Publish Date	2020-12-29
Publish Year	2020
Doi	DOI: 10.15294/biosaintifika.v12i3.23482
Citation	
Source	Biosaintifika: Journal of Biology & Biology Education
Source Issue	Vol 12, No 3 (2020): December 2020
Source Page	453-458
Url	https://journal.unnes.ac.id/nju/index.php/biosaintifika/article/view/23482/11193
Author	SUGIYONO, S.Si, Ph.D