## The Effectiveness of SCIDOTS to Reduce Smoking Behavior Among Patients with Pulmonary Tuberculosis : Systematic Review

The Effectiveness of SCIDOTS to Reduce Smoking Behavior Among Patients with Pulmonary Tuberculosis: Systematic Review
2 of 3
4
Aims: The aim of this study is to identify and summarize the results of previous studies on the use of SCIDOTS to reduce smoking cessation in patients with pulmonary tuberculosis. Methods: The research used was a literature review of three articles. The assessment of the articles shall be measured using the CASP (Critical Appraisal Skills Program) format. Results: The result of this study incorporating smoking cessation programs into pulmonary TB control programs, as new diagnoses can increase cessation rates by 71.7%. Various interventions, including counseling sessions, behavioral therapy, and health promotion programs, have proven successful. Conclusions: The study found that the intervention SCIDOTS, or quitting smoking, is highly effective in treating tuberculosis patients, accelerating their active treatment.
STIKep PPNI Jawa Barat
2024-08-31
2024
DOI: 10.33755/jkk.v10i4.700
Jurnal Keperawatan Komprehensif (Comprehensive Nursing Journal)
Vol. 10 No. 4 (2024): JURNAL KEPERAWATAN KOMPREHENSIF (COMPREHENSIVE NURSING JOURNAL) : SPECIAL EDIT
425-434
http://journal.stikep-ppnijabar.ac.id/index.php/jkk/article/view/700/467
Dr NUNUNG NURHAYATI, S.Si, M.Si