

PROFIL ASAM LEMAK RANTAI CABANG CAIRAN RUMEN YANG DISUPLEMENTASI LEGUMINOSA DALAM RANSUM BERBASIS INDEKS SINKRONISASI PROTEIN-ENERGI

Title	PROFIL ASAM LEMAK RANTAI CABANG CAIRAN RUMEN YANG DISUPLEMENTASI LEGUMINOSA DALAM RANSUM BERBASIS INDEKS SINKRONISASI PROTEIN-ENERGI
Author Order	of
Accreditation	
Abstract	<p>The experiment was aimed to assess the use of the legume and synchronization index of protein and energy (SPE) in the diet of dairy cattles on branch chain volatile fatty acids (BCVFA) profile. The research was applied in In vitro technique. The research was used a Completely Randomized Design (CRD), factorially pattern (2x3), the first factor was the two species of legume (Sesbania and Leucaena) and the second factor was the three levels of the SPE index (0.4, 0.5, and 0.6), there were 6 treatments combination and replicates 4 times each. The results showed interaction between legumes with SPE index was highly significant ($P < 0.01$) upon iso butirat, but was nonsignificant ($P > 0.05$) on iso valerat and valerat. Legume and SPE index respectively showed nonsignificant ($P > 0.05$) on iso valerat and valerat. It can be concluded that Turi supplementation with high SPE index (0.6) gave the best effect on Branch Chain Volatile Fatty Acids (BCVFA) profile. $\hat{\Delta}$ Key words: Branch chain volatile fatty acids, Protein-energy synchronization index, Legume</p> <p>The experiment was aimed to assess the use of the legume and synchronization index of protein and energy (SPE) in the diet of dairy cattles on branch chain volatile fatty acids (BCVFA) profile. The research was applied in In vitro technique. The research was used a Completely Randomized Design (CRD), factorially pattern (2x3), the first factor was the two species of legume (Sesbania and Leucaena) and the second factor was the three levels of the SPE index (0.4, 0.5, and 0.6), there were 6 treatments combination and replicates 4 times each. The results showed interaction between legumes with SPE index was highly significant ($P < 0.01$) upon iso butirat, but was nonsignificant ($P > 0.05$) on iso valerat and valerat. Legume and SPE index respectively showed nonsignificant ($P > 0.05$) on iso valerat and valerat. It can be concluded that Turi supplementation with high SPE index (0.6) gave the best effect on Branch Chain Volatile Fatty Acids (BCVFA) profile. $\hat{\Delta}$ Key words: Branch chain volatile fatty acids, Protein-energy synchronization index, Legume</p>
Publisher Name	Universitas Tidar
Publish Date	2018-03-28
Publish Year	2018
Doi	
Citation	
Source	Journal of Livestock Science and Production
Source Issue	Vol 2, No 1 (2018): Journal of Livestock Science and Production
Source Page	50-55
Url	http://jurnal.untidar.ac.id/index.php/jalspro/article/view/683
Author	AFDUHA NURUS SYAMSI, S.Pt, M.P