

In vitro study of DNA adduct 8-hidroxy-2'-deoxyguanosine (8-OHdG) formation through Fenton-like reaction with butylated hydroxytoluene quinone (BHT quinone)

<b>Title</b>	In vitro study of DNA adduct 8-hidroxy-2'-deoxyguanosine (8-OHdG) formation through Fenton-like reaction with butylated hydroxytoluene quinone (BHT quinone)
<b>Abstract</b>	
<b>Authors</b>	B Budiawan, S Handayani, IC Dani, R Bakri, S Hannaa, F Irawati
<b>Journal Name</b>	AIP Conference Proceedings 2023 (1), 2018
<b>Publish Year</b>	2018
<b>Citation</b>	(not set)
<b>Url</b>	<a (8-ohdg)="" (bht="" 8-hidroxy-2'-deoxyguanosine="" adduct="" butylated="" dna="" fenton-like="" formation="" href="https://scholar.google.com/scholar?q=+intitle:" hydroxytoluene="" in="" of="" quinone="" quinone)\""="" reaction="" study="" through="" vitro="" with="">https://scholar.google.com/scholar?q=+intitle:"In vitro study of DNA adduct 8-hidroxy-2'-deoxyguanosine (8-OHdG) formation through Fenton-like reaction with butylated hydroxytoluene quinone (BHT quinone)"</a>
<b>Author</b>	Dr SANTI NUR HANDAYANI, S.Si, M.Si