

Potential of Various Rice-Washing Water as a Source of Manufacturing Secondary Metabolites of Trichoderma harzianum T10 to Control Cucumber Crown and Root Rot

<b>Title</b>	Potential of Various Rice-Washing Water as a Source of Manufacturing Secondary Metabolites of Trichoderma harzianum T10 to Control Cucumber Crown and Root Rot
<b>Abstract</b>	
<b>Authors</b>	L Soesanto, MA Dapawole, MWR Sastyawan, E Mugiastuti, A Manan, ...
<b>Journal Name</b>	Agrotechnology Research Journal 8 (2), 0
<b>Publish Year</b>	(not set)
<b>Citation</b>	(not set)
<b>Url</b>	<a a="" and="" as="" control="" crown="" cucumber="" harzianum="" href="https://scholar.google.com/scholar?q=+intitle:" manufacturing="" metabolites="" of="" potential="" rice-washing="" root="" rot"="" secondary="" source="" t10="" to="" trichoderma="" various="" water="">https://scholar.google.com/scholar?q=+intitle:"Potential of Various Rice-Washing Water as a Source of Manufacturing Secondary Metabolites of Trichoderma harzianum T10 to Control Cucumber Crown and Root Rot"</a>
<b>Author</b>	Dr ENDANG MUGIASTUTI, S.P, M.P