## Incidence of malaria is clustered and buffers around plantations: a spatial analysis

Publons ID	32441357
Wos ID	WOS:000376161700009
Doi	10.18051/UnivMed.2015.v34.138-148
Title	Incidence of malaria is clustered and buffers around plantations: a spatial analysis
First Author	Wibowo, Yudhi; Laksana, Agung Saprasetya Dwi; Mulyanto, Joko;
Last Author	Purnomo, Agus Y.
Authors	Wibowo, Y; Laksana, ASD; Mulyanto, J; Wicaksono, MA; Purnomo, AY;
Publish Date	MAY-AUG 2015
Journal Name	UNIVERSA MEDICINA
Citation	
Abstract	a:8:{i:0;s:10:"BACKGROUND";i:1;s:437:"Malaria is re-emerging because of imported cases and the presence of potential vectors that can transmit and spread malaria. Malaria is a health problem in Banyumas District. Mapping the spread of infectious diseases is epidemiologically important. The purpose of this study was to determine the relationship between the variables and the epidemiology of malaria that were spatially modeled using the geographic information system (GIS).";i:2;s:7:"METHODS";i:3;s:385:"This was a case-control study with ratio of 1:1. Cases were malaria-positive patients and controls were people without malaria, as diagnosed by microscopic examination. Minimum sample size was 139 per group and total sample size was 282 people. Chi-square was used to test the relationship between the variables, and GIS modeling to determine the spatial distribution of malaria cases.";i:4;s:7:"RESULTS";i:5;s:506:"There were significant relationships between level of income below minimum wage, not using mosquito nets, not using wire netting, not using insect repellents, habit of going out at night, history of malaria, cattle sheds not located between woods and residential area, history of going to endemic areas, residence at distances <1000 m from plantations, bushes, swamps and puddles, with incidence of confirmed malaria (p<0.001). The group of cases living <= 1000 meters from plantations numbered 141 (100%).";i:6;s:11:"CONCLUSIONS";i:7;s:229:"Malaria incidence is clustered and buffers around plantations at <= 1000 m. Malaria hot spots are displayed as risk maps that are useful for monitoring and spatial targeting of prevention and control measures against the disease.";}
Publish Type	Journal
Publish Year	2015
Page Begin	138
Page End	148
lssn	1907-3062
Eissn	2407-2230
Url	https://www.webofscience.com/wos/woscc/full-record/WOS:000376161700009
Author	dr. JOKO MULYANTO, S.Ked, M.Sc., PhD